

NERIDRONIC ACID FOR TREATING COMPLEX REGIONAL PAIN SYNDROME

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Prov. App. No. 62/431,287, filed Dec. 7, 2016; this application is also a continuation-in-part of U.S. patent application Ser. No. 15/357,932, filed Nov. 21, 2016; which is a continuation-in-part of U.S. patent application Ser. No. 14/530,556, filed Oct. 31, 2014, which is a continuation-in-part of U.S. patent application Ser. No. 14/279,229, filed May 15, 2014, now U.S. Pat. No. 9,034,889, which is a continuation of U.S. patent application Ser. No. 14/063,979, filed Oct. 25, 2013, now U.S. Pat. No. 8,802,658, which is a continuation-in-part of U.S. patent application Ser. No. 13/894,274, filed May 14, 2013, now abandoned, which claims the benefit of U.S. Prov. App. No. 61/646,538, filed May 14, 2012; the above U.S. patent application Ser. No. 15/357,932 is also a continuation-in-part of International Pat. App. No. PCT/US2015/032739, filed May 27, 2015, which is a continuation of International Pat. App. No. PCT/US2014/050427, filed Aug. 8, 2014, which is a continuation of U.S. patent application Ser. No. 14/279,241, filed May 15, 2014, now abandoned; any of the applications, U.S. patents issued from, or U.S. publications of any of the above applications are incorporated by reference in their entirety.

SUMMARY

Bisphosphonate compounds are potent inhibitors of osteoclast activity, and are used clinically to treat bone-related conditions such as osteoporosis and Paget's disease of bone; and cancer-related conditions including multiple myeloma, and bone metastases from solid tumors. They generally have low oral bioavailability.

Patchy osteoporosis and bone marrow edema may result from osteoclast hyperactivity. Zoledronic acid is a potent inhibitor of bone resorption and osteoclast activity. Nitrogen containing bisphosphonates, such as zoledronic acid, also inhibit the mevalonate pathway in the osteoclast thereby interrupting normal osteoclast function.

It has been discovered that oral dosage forms of bisphosphonate compounds, such as zoledronic acid, can be used to treat or alleviate pain or related conditions.

Some embodiments include a method of enhancing the oral bioavailability of zoledronic acid comprising orally administering a dosage form containing zoledronic acid in the disodium salt form.

Some embodiments include a dosage form comprising zoledronic acid in the disodium salt form, wherein the bioavailability, in a mammal, of zoledronic acid in the disodium salt form is greater than the bioavailability of zoledronic acid in the diacid form would be in the same dosage form.

Some embodiments include a dosage form comprising zoledronic acid in an acid or a salt form, such as the disodium salt form, wherein the dosage form contains an amount of zoledronic acid in the disodium salt form that provides an area under the plasma concentration curve of zoledronic acid of about 4 ng·h/mL to about 2000 ng·h/mL to a human being to which the dosage form is administered.

Some embodiments include a dosage form comprising zoledronic acid in the disodium salt form, wherein the disodium salt form is present in a lower molar amount than would be present if the zoledronic acid were in the diacid form; and wherein the zoledronic acid in the disodium salt form has an improved bioavailability as compared to the zoledronic acid in the diacid form to the extent that the lower

molar amount of the disodium salt in the dosage form does not reduce the amount of zoledronic acid delivered to the plasma of a mammal.

Although an oral dosage form with enhanced bioavailability with respect to the bisphosphonate compound can be used, the treatment can also be effective using an oral dosage form that includes a bisphosphonate compound, such as zoledronic acid, wherein the bioavailability of the bisphosphonate is unenhanced, or is substantially unenhanced.

Some embodiments include a method of relieving inflammatory pain comprising administering an oral dosage form containing zoledronic acid to a mammal in need thereof, wherein the mammal experiences significant pain relief more than 3 hours after administration of the dosage form.

Some embodiments include a method of relieving pain associated with an arthritis comprising administering an oral dosage form containing zoledronic acid to a human being in need thereof.

Some embodiments include a method of treating complex regional pain syndrome comprising administering an oral dosage form containing zoledronic acid to a mammal in need thereof.

Some embodiments include an oral dosage form comprising zoledronic acid, wherein the oral bioavailability of zoledronic acid is substantially unenhanced. For example, in some embodiments, the oral bioavailability in the dosage form is about 0.01% to about 4%.

Some embodiments include a pharmaceutical product comprising more than one unit of an oral dosage form described herein. In some embodiments, each unit of the oral dosage form contains about 1 mg to about 50 mg of zoledronic acid.

Some embodiments include a method of relieving inflammatory pain comprising administering an oral dosage form containing zoledronic acid to a mammal in need thereof.

In some embodiments, the mammal receives a total monthly dose of zoledronic acid that is about 800 mg/m² or less.

In some embodiments, the dosage form contains about 10 mg/m² to about 20 mg/m² based upon the body surface area of the mammal.

Some embodiments include a method of relieving inflammatory pain comprising orally administering zoledronic acid to a mammal in need thereof.

In some embodiments, about 300 mg/m² to about 600 mg/m² of zoledronic acid is administered per month, based upon the body surface area of the mammal.

In some embodiments, about 50 mg/m² to about 600 mg/m² of zoledronic acid is administered per month, based upon the body surface area of the mammal.

Some embodiments include administering an osteoclast inhibitor, such as a bisphosphonate, including zoledronic acid, neridronic acid, etc. to inhibit the development of pain, unweighting, and edema when administered early such as when a precipitating event such as fracture occurs, wherein the precipitating event is associated with CRPS.

Some embodiments include administering an osteoclast inhibitor, such as a bisphosphonate, including zoledronic acid, neridronic acid, etc. to reverse established allodynia and unweighting when administered at least 4 weeks after a precipitating event such as fracture that is associated with CRPS.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 is a plot of pain compression thresholds in a rat model of inflammatory pain using three different doses of zoledronic acid. Measurements were taken at baseline (BL) and at various time points after dosing on the days indicated.